
Hesi Case Study Pediatric Congenital Heart Disease

*Hesi Case
Study
Pediatric
Congenital
Heart
Disease*

*Downloaded from
www2.lacavedespapilles.com
by Danika & Carmelo*

UNDERSTANDING HESI CASE STUDY PEDIATRIC CONGENITAL HEART DISEASE: A COMPREHENSIVE GUIDE FOR NURSING STUDENTS

The HESI exam is a pivotal milestone for nursing students, and few topics evoke as much anxiety as pediatric congenital heart disease. Mastering the **HESI**

case study pediatric congenital heart disease requires not only memorizing cardiac defects but also understanding the pathophysiology, nursing interventions, and family-centered care strategies. This article provides a deep dive into the key concepts, common defects, and practical approaches to help you excel in your HESI pediatric case study.

What

Makes

the HESI

Case

Study

Pediatric

Congenital

Heart

Disease

Unique?

The HESI pediatric case study on congenital heart disease is designed to test your critical thinking and clinical judgment. Unlike straightforward multiple-choice questions, the case study presents a

realistic patient scenario that evolves over time. You must analyze symptoms, interpret diagnostic data, prioritize nursing interventions, and anticipate complications. The **HESI case study pediatric congenital heart disease** often includes a child with a cyanotic or acyanotic defect, requiring you to differentiate between the two and tailor your care accordingly.

Common defects featured in these case studies include ventricular septal defect (VSD), tetralogy of Fallot (TOF), transposition of the great arteries (TGA), and patent ductus arteriosus (PDA). Each defect has distinct hemodynamic consequences, and the HESI expects you to

connect the dots between anatomy, physiology, and clinical presentation.

Key Concepts to Master for the HESI Pediatric Congenit al Heart Disease Case

Study

To succeed in the **HESI case study pediatric congenital heart disease**, you need a solid grasp of several foundational concepts. These include fetal circulation, the transition to neonatal circulation, and the classification of congenital heart defects.

FETAL VS. NEONATAL CIRCULATION

Understanding fetal circulation is critical because many congenital heart defects become symptomatic when the ductus arteriosus closes after birth. In fetal circulation, the ductus arteriosus shunts blood from the pulmonary artery to

the aorta, bypassing the non-functioning lungs. After birth, this structure normally closes within days. However, in defects like TGA or TOF, the infant may depend on the ductus for survival, and its closure can precipitate a life-threatening crisis. The **HESI case study pediatric congenital heart disease** often tests your ability to recognize when prostaglandin E1 infusion is needed to keep the ductus open.

CLASSIFICATION OF CONGENITAL HEART DEFECTS

Congenital heart defects are broadly classified as acyanotic or cyanotic. Acyanotic defects, such as VSD and PDA, involve left-to-right shunting that

increases pulmonary blood flow. Cyanotic defects, such as TOF and TGA, involve right-to-left shunting that sends deoxygenated blood into the systemic circulation. The HESI case study will challenge you to identify the defect type based on oxygenation levels, heart murmurs, and chest X-ray findings.

- **Acyanotic defects:**

Increased pulmonary blood flow, normal oxygen saturation initially, risk of heart failure.

- **Cyanotic defects:**

Decreased pulmonary blood flow or mixed circulation, low oxygen saturation,

cyanosis.

- **Obstructive defects:**
Coarctation of the aorta, aortic stenosis, pulmonary stenosis—these can cause pressure overload.

pediatric congenital heart disease.

Understanding their pathophysiology, clinical features, and nursing management is essential.

VENTRICULAR SEPTAL DEFECT (VSD)

Common Congenital Heart Defects in HESI Case Studies

Let us explore the most frequently encountered defects in the **HESI case study**

VSD is the most common congenital heart defect. It involves an opening in the septum between the ventricles, causing left-to-right shunting. This increases pulmonary blood flow and can lead to heart failure. On assessment, you may hear a holosystolic murmur at the left lower sternal border. Infants with a large VSD often present with tachypnea, poor feeding, and failure to thrive. Surgical repair

is usually scheduled before school age, but small VSDs may close spontaneously.

In the **HESI case study pediatric congenital heart disease**, you might be asked to prioritize interventions such as administering digoxin and furosemide for heart failure, providing high-calorie formula, and positioning the infant in a semi-upright position to ease breathing. Be prepared to recognize signs of pulmonary hypertension, which can reverse the shunt and cause Eisenmenger syndrome.

TETRALOGY OF FALLOT (TOF)

TOF is the most common cyanotic heart defect and a frequent

topic in HESI case studies. It comprises four anatomical abnormalities:

ventricular septal defect, overriding aorta, right ventricular hypertrophy, and pulmonary stenosis. The severity of pulmonary stenosis determines the degree of cyanosis. Infants with TOF may experience "tet spells," which are hypercyanotic episodes triggered by crying or feeding. During a tet spell, the infant becomes deeply cyanotic, tachypneic, and irritable.

The **HESI case study pediatric congenital heart disease** will test your ability to manage a tet spell. Interventions include placing the infant in the knee-chest position to increase systemic

vascular resistance, administering oxygen, morphine to reduce agitation, and intravenous fluids. Long-term management involves surgical correction, typically performed in the first year of life. You should also understand that these infants are at risk for infective endocarditis and may require prophylactic antibiotics before dental procedures.

TRANSPOSITION OF THE GREAT ARTERIES (TGA)

TGA is a cyanotic defect where the aorta arises from the right ventricle and the pulmonary artery from the left ventricle, creating two parallel circulations. This defect is incompatible

with life unless there is mixing of blood through a patent ductus arteriosus, atrial septal defect, or VSD. Newborns present with severe cyanosis shortly after birth. The immediate intervention is to start a prostaglandin E1 infusion to maintain ductal patency, followed by a balloon atrial septostomy (Rashkind procedure) to improve mixing.

In the **HESI case study pediatric congenital heart disease**, you may need to prioritize the administration of prostaglandin E1 and prepare the infant for surgery. The definitive treatment is the arterial switch operation, which should be performed in the first two weeks of life. Postoperative care

involves monitoring for arrhythmias, low cardiac output, and bleeding.

PATENT DUCTUS ARTERIOSUS (PDA)

PDA is an acyanotic defect where the ductus arteriosus fails to close after birth. This creates a left-to-right shunt from the aorta to the pulmonary artery, increasing pulmonary blood flow. On assessment, you may hear a continuous "machinery" murmur. Preterm infants are at higher risk for PDA, and it can exacerbate respiratory distress syndrome.

The **HESI case study pediatric congenital heart disease** might present a preterm infant with PDA. Management includes fluid restriction,

diuretics, and indomethacin or ibuprofen to promote ductal closure. If medical management fails, surgical ligation may be necessary. You should also monitor for signs of heart failure and feed the infant with high-calorie formula to support growth.

Nursing Assessment and Diagnostic Tools

A thorough assessment is the cornerstone of managing congenital heart disease. In the **HESI case study pediatric congenital heart disease**, you

will be expected to interpret vital signs, oxygen saturation, and diagnostic tests.

PHYSICAL ASSESSMENT FINDINGS

Key assessment findings include cyanosis, clubbing of the fingers and toes, tachypnea, tachycardia, and poor weight gain. Auscultate for murmurs, gallops, and rubs. Palpate for hepatomegaly, which indicates right-sided heart failure. In infants, observe for diaphoresis during feeding, a sign of increased work of breathing. The **HESI case study pediatric congenital heart disease** often includes a description of the infant's appearance, and you must correlate these findings with the

specific defect.

DIAGNOSTIC TESTS AND PROCEDURES

Common diagnostic tests include:

- **Chest X-ray:** May show cardiomegaly, increased pulmonary vascular markings (acyanotic defects), or a boot-shaped heart (TOF).
- **Echocardiogram:** The gold standard for diagnosing structural defects. It visualizes the anatomy and blood flow.
- **Electrocardiogram (ECG):** Can show chamber hypertrophy and axis deviations.

- **Cardiac catheterization**

: Used to measure pressures, oxygen saturations, and sometimes to perform interventions like balloon septostomy.

- **Pulse oximetry:** Screening tool for critical congenital heart disease in newborns. Low preductal and postductal saturations suggest a ductal-dependent lesion.

In the **HESI case study pediatric congenital heart disease**, you may be given results from these tests and asked to interpret them. For example, a boot-

shaped heart on X-ray with a history of tet spells points strongly to TOF.

Nursing Interventions and Family-Centered Care

Caring for a child with congenital heart disease extends beyond medical management. The **HESI case study pediatric congenital heart disease** emphasizes holistic care that includes the family.

PREOPERATIVE AND POSTOPERATIVE CARE

Preoperative care focuses on stabilizing the infant. This may involve administering medications such as digoxin to improve contractility, diuretics to reduce fluid overload, and oxygen therapy. Maintain adequate nutrition by using high-calorie formulas or nasogastric feedings, as these infants tire easily during feeding. Monitor for signs of infection and provide prophylactic antibiotics as ordered.

Postoperative care after cardiac surgery requires close monitoring in the pediatric intensive care unit. Assess vital signs frequently, monitor chest tube output, and

observe for bleeding. Pain management is crucial because pain increases oxygen consumption. The **HESI case study pediatric congenital heart disease** may test your ability to prioritize assessments, such as checking for decreased urine output, which could indicate low cardiac output syndrome.

PARENT EDUCATION AND PSYCHOSOCIAL SUPPORT

Parents of children with congenital heart disease face immense stress. They need clear explanations about the defect, treatment plan, and home care. Teach them to recognize signs of worsening heart failure, such as increased respiratory rate, decreased

appetite, or irritability. Instruct them on medication administration, wound care, and when to seek emergency care.

In the **HESI case study pediatric congenital heart disease**, you may encounter a scenario where the parents are anxious or have misconceptions. Your response should include active listening, providing factual information in simple terms, and offering emotional support. Refer them to support groups and resources such as the American Heart Association. Remember that family-centered care improves outcomes and adherence to treatment.

Pharmacology and Medication Management

Medications play a key role in managing congenital heart disease, and the **HESI case study pediatric congenital heart disease** will test your knowledge of commonly used drugs.

- **Digoxin:**
Increases myocardial contractility and slows heart rate. Monitor for toxicity, which can cause

bradycardia, nausea, and visual disturbances. Check apical pulse for one minute before administration; hold if heart rate is below 90-110 bpm (depending on age).

- **Furosemide:** A loop diuretic that reduces fluid overload. Monitor electrolytes, especially potassium, and watch for ototoxicity. Weigh the infant daily to assess fluid status.
- **Captopril:** An ACE inhibitor that reduces afterload and improves cardiac output. Monitor blood pressure and renal function.

- **Prostaglandin E1:** Used to maintain ductal patency in ductal-dependent lesions. Monitor for apnea, hypotension, and fever. Have resuscitation equipment available.
- **Indomethacin or Ibuprofen:** Used to close PDA in preterm infants. Monitor renal function and platelet count.

Understanding the indications, side effects, and nursing considerations for these medications is essential for the HESI case study.

Common Complications and Emergency Situations

The **HESI case study pediatric congenital heart disease** often includes complications that require immediate intervention. Being prepared for these emergencies is critical.

HEART FAILURE IN INFANTS

Heart failure is a common complication of acyanotic defects with large left-to-right

shunts. Signs include tachypnea, tachycardia, poor feeding, diaphoresis, and hepatomegaly. Management involves oxygen, digoxin, diuretics, and afterload reduction. In severe cases, mechanical ventilation or inotropic support may be needed. The HESI case study may ask you to prioritize interventions, such as administering oxygen first before other treatments.

HYPERCYANOTIC (TET) SPELLS

As discussed earlier, tet spells occur in TOF. The immediate priority is to place the infant in the knee-chest position to increase systemic vascular resistance and decrease the right-to-left shunt.